

Some observations on a Kapteyn series

Diego Dominici *

Department of Mathematics
State University of New York at New Paltz
75 S. Manheim Blvd. Suite 9
New Paltz, NY 12561-2443
USA

Phone: (845) 257-2607

Fax: (845) 257-3571

Abstract

We study the Kapteyn series $\sum_{n=1}^{\infty} t^n J_n(nz)$. We find a series representation in powers of z and analyze its radius of convergence.

Keywords: Kapteyn series, power series, Bessel functions.

MSC-class: 42C10 (Primary) 30B50, 33C10 (Secondary).

1 Introduction

Series of the form

$$\sum_{n=1}^{\infty} \alpha_n J_{n+\nu}[(n+\nu)z], \quad \nu \in \mathbb{C} \quad (1)$$

where $J_n(\cdot)$ is the Bessel function of the first kind, are called *Kapteyn series*. The series (1) is convergent and represents an analytic function throughout the domain in which [8, 17.3]

$$\Omega(z) < \lim_{n \rightarrow \infty} |\alpha_n|^{-(\nu+n)}, \quad (2)$$

*e-mail: dominicd@newpaltz.edu

where

$$\Omega(z) = \left| \frac{z \exp [\sqrt{1-z^2}]}{1 + \sqrt{1-z^2}} \right|. \quad (3)$$

Kapteyn series first appeared in Lagrange's solution [7]

$$E = M + 2 \sum_{n=1}^{\infty} \frac{1}{n} J_n(\varepsilon n) \sin(Mn) \quad (4)$$

of Kepler's equation [3]

$$M = E - \varepsilon \sin(M).$$

The solution (4) was independently discovered by Friedrich Bessel in [2], where he introduced the functions which now bear his name.

Kapteyn series were systematically studied by Willem Kapteyn in his article [5], where he proved the following expansion theorem.

Theorem 1 *Let $f(z)$ be a function which is analytic throughout the region*

$$D_a = \{z \in \mathbb{C} \mid \Omega(z) \leq a\},$$

with $a \leq 1$. Then,

$$f(z) = \alpha_0 + 2 \sum_{n=1}^{\infty} \alpha_n J_n(nz), \quad z \in D_a$$

where

$$\alpha_n = \frac{1}{2\pi i} \oint \Theta_n(z) f(z) dz$$

and the path of integration is the curve on which $\Omega(z) = a$. The function $\Theta_n(z)$ is the Kapteyn polynomial defined by

$$\begin{aligned} \Theta_0(z) &= \frac{1}{z} \\ \Theta_n(z) &= \frac{1}{4} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(n-2k)^2 (n-k-1)!}{k!} \left(\frac{nz}{2}\right)^{2k-n}, \quad n \geq 1, \end{aligned}$$

where $\lfloor \cdot \rfloor$ denotes the integer-part function.

Proof. See [8, 17.4]. ■

As a corollary [8], one finds that if the Taylor series for $f(z)$ is

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

then,

$$\begin{aligned} \alpha_0 &= a_0 \\ \alpha_n &= \frac{1}{4} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(n-2k)^2 (n-k-1)!}{k! \left(\frac{n}{2}\right)^{n-2k+1}} a_{n-2k}, \quad n \geq 1. \end{aligned} \quad (5)$$

In [6], Kapteyn established the fundamental formula

$$\frac{1}{1-z} = 1 + 2 \sum_{n=1}^{\infty} J_n(nz), \quad \Omega(z) < 1, \quad (6)$$

using the integral representation [8]

$$J_n(nz) = \frac{1}{2\pi i} \int_{\mathcal{H}} \left\{ \frac{\exp\left[\frac{z}{2}\left(s - \frac{1}{s}\right)\right]}{s} \right\}^n \frac{ds}{s},$$

where the Hankel contour $\mathcal{H}$ encircles the origin once counterclockwise.

The purpose of this paper is to generalize (6) by means of the function

$$F(z, t) = \sum_{n=1}^{\infty} t^n J_n(nz), \quad z \in \mathbb{C}, \quad t \in \mathbb{R}, \quad (7)$$

so that (6) corresponds to the particular case $t = 1$. In Section 2 we write $F(z, t)$ as a series in powers of z

$$F(z, t) = \sum_{n=1}^{\infty} A_n(t) z^n \quad (8)$$

and find the coefficients $A_n(t)$. Although one could replace $\alpha_n = t^n$ in (5) and solve the problem

$$t^n = \frac{1}{4} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(n-2k)^2 (n-k-1)!}{k! \left(\frac{n}{2}\right)^{n-2k+1}} A_{n-2k}(t),$$

we take a different approach that only uses the differential equation satisfied by the Bessel functions. In Section 3 we analyze the radius of convergence of (8) for different ranges of t . Finally, in Section 4 we solve the inversion problem (5) for arbitrary α_n .

2 Power series

Since the Bessel function $J_n(z)$ is a solution of [8]

$$z^2 J_n'' + z J_n' + (z^2 - n^2) J_n = 0,$$

the function $y_n(z) = J_n(nz)$ satisfies the ODE

$$z^2 y_n'' + z y_n' + n^2 (z^2 - 1) y_n = 0. \quad (9)$$

We also observe that

$$t^2 (t^n)'' + t (t^n)' = n^2 t^n. \quad (10)$$

Using (9) and (10) in (7), we have

$$z^2 F_{zz} + z F_z = (1 - z^2) (t^2 F_{tt} + t F_t), \quad (11)$$

where the subscripts denote partial derivatives.

Replacing (8) in (11), we get the equation

$$t^2 A_n'' + t A_n' - n^2 A_n = t^2 A_{n-2}'' + t A_{n-2}', \quad n \geq 1, \quad (12)$$

where we define $A_n \equiv 0$ for $n = -1, 0$.

From (7) and (8), we have

$$0 = F(z, 0) = \sum_{n=1}^{\infty} A_n(0) z^n$$

and, from (6), we see that

$$\sum_{n=1}^{\infty} \frac{1}{2} z^n = \frac{z}{2(1-z)} = F(z, 1) = \sum_{n=1}^{\infty} A_n(1) z^n,$$

which together imply that

$$A_n(0) = 0, \quad A_n(1) = \frac{1}{2}, \quad n \geq 1. \quad (13)$$

Solving (12) with the boundary conditions (13), we obtain

$$\begin{aligned} A_1(t) &= \frac{t}{2}, & A_2(t) &= \frac{t^2}{2}, & A_3(t) &= \frac{9}{16}t^3 - \frac{1}{16}t \\ A_4(t) &= \frac{2}{3}t^4 - \frac{1}{6}t^2, & A_5(t) &= \frac{625}{768}t^5 - \frac{81}{256}t^3 + \frac{1}{384}t, \dots \end{aligned}$$

Thus, the function $A_n(t)$ is a polynomial in t of degree n . Writing

$$A_n(t) = \sum_{k=0}^n C_k^n t^k \quad (14)$$

and using (12), we get the recurrence

$$\begin{aligned} (n^2 - k^2) C_k^n &= -k^2 C_k^{n-2}, & \text{for } k = 0, 1, \dots, n-2 \\ (n^2 - k^2) C_k^n &= 0, & \text{for } k = n-1, n. \end{aligned} \quad (15)$$

To solve (15), we recall that the double factorial function satisfies

$$(n-k)!! = (n-k)(n-k-2)!!, \quad (n+k)!! = (n+k)(n+k-2)!!,$$

which gives

$$(n^2 - k^2) \frac{1}{(n-k)!! \times (n+k)!!} = \frac{1}{(n-k-2)!! \times (n+k-2)!!}.$$

This suggests the solution

$$C_k^n = \frac{\cos\left(\frac{n-k}{2}\pi\right) k^n}{(n-k)!! \times (n+k)!!}, \quad 0 \leq k \leq n. \quad (16)$$

Therefore,

$$A_n(t) = \sum_{k=0}^n \frac{\cos\left(\frac{n-k}{2}\pi\right) k^n}{(n-k)!! \times (n+k)!!} t^k \quad (17)$$

or, after rearranging terms,

$$A_n(t) = \frac{(-1)^n}{n!} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n}{k} \left(k - \frac{n}{2}\right)^n t^{n-2k}, \quad (18)$$

It is clear from (17) that $A_n(0) = 0$. To show that $A_n(1) = \frac{1}{2}$, we use the identity [4, 0.154, #5]

$$\sum_{k=0}^n (-1)^k \binom{n}{k} (k + \alpha)^n = (-1)^n n!, \quad (19)$$

where α is arbitrary. From (18) we have

$$A_n(t) + A_n\left(\frac{1}{t}\right) = \frac{(-1)^n}{n!} \sum_{k=0}^n (-1)^k \binom{n}{k} \left(k - \frac{n}{2}\right)^n t^{n-2k}. \quad (20)$$

Setting $t = 1$ in (20), we have

$$2A_n(1) = \frac{(-1)^n}{n!} \sum_{k=0}^n (-1)^k \binom{n}{k} \left(k - \frac{n}{2}\right)^n$$

and from (19), we conclude that $2A_n(1) = 1$.

3 Radius of convergence

We shall now find the radius of convergence $R(t)$ of the power series (8). From (7) we observe that $R(t) \rightarrow 0$ as $t \rightarrow \infty$ and $R(t) \rightarrow \infty$ as $t \rightarrow 0$, while (6) gives $R(1) = 1$ (see Figure 1). Since from (18) we have $A_n(-t) = (-1)^n A_n(t)$, we shall limit our analysis to $t > 0$.

When $t \gg 1$, the largest term in the sum (18) corresponds to $k = 0$, and therefore

$$A_n(t) \sim \frac{1}{n!} \left(\frac{nt}{2}\right)^n, \quad t \rightarrow \infty. \quad (21)$$

Using Stirling's formula [1]

$$n! \sim \sqrt{2\pi n} n^n e^{-n}, \quad n \rightarrow \infty$$

we obtain

$$A_n(t) \sim \frac{1}{\sqrt{2\pi n}} \left(\frac{et}{2}\right)^n, \quad t \rightarrow \infty. \quad (22)$$

Thus,

$$R(t) = \lim_{n \rightarrow \infty} |A_n(t)|^{-\frac{1}{n}} \sim \frac{2e^{-1}}{t}, \quad t \rightarrow \infty. \quad (23)$$




Figure 1: A sketch of $|A_{500}(t)|^{-\frac{1}{500}} \simeq R(t)$.




Figure 2: A sketch of $\ln [|A_n(0.1)|]$.

When $t < 1$, (18) is highly oscillatory and there is no approximation like (21) valid in this range.

To find a formula for $R(t)$ valid for all t , we observe that, for a fixed value $t > 0$, we have

$$\ln [|A_n(t)|] \simeq \beta(t)n, \quad (24)$$

for some function $\beta(t)$ (see Figure 2).

Thus, we consider an asymptotic expansion of the form

$$A_n(t) \sim \frac{1}{2}\theta_n(t) [\psi(t)]^n, \quad n \rightarrow \infty, \quad (25)$$

where $\theta_n(t)$ denotes the sign of $A_n(t)$ and $\psi(1) = 1$. Replacing (25) in (12) and setting

$$\theta'_n(t) = \theta''_n(t) = 0, \quad \theta'_{n-2}(t) = \theta''_{n-2}(t) = 0,$$

we obtain, to leading order in n ,

$$(t\psi')^2 (\theta_n \psi^2 - \theta_{n-2}) - \theta_n \psi^4 = 0. \quad (26)$$

Since

$$\theta_n = \pm 1, \quad \theta_{n-2} = \pm 1,$$

and $\psi(t)$ is increasing for $t > 0$, we get

$$\psi' = \frac{\psi^2}{t\sqrt{\psi^2 \pm 1}}. \quad (27)$$

Solving (27) subject to $\psi(1) = 1$, we obtain the implicit solutions

$$\ln(t) + \frac{[\psi^2(t) + 1]^{\frac{3}{2}}}{\psi(t)} - \psi(t)\sqrt{\psi^2(t) + 1} - \operatorname{arcsinh}[\psi(t)] - \sqrt{2} + \ln(\sqrt{2} + 1) = 0, \quad (28)$$

for $0 < t \leq 1$ and

$$\ln(t) - \frac{[\psi^2(t) - 1]^{\frac{3}{2}}}{\psi(t)} + \psi(t)\sqrt{\psi^2(t) - 1} - \ln[\psi(t) + \sqrt{\psi^2(t) - 1}] = 0 \quad (29)$$

for $t \geq 1$.

Although we cannot solve (28) and (29) exactly, we can consider the limiting cases as $t \rightarrow 0$ and $t \rightarrow \infty$. Since $\psi(t) \rightarrow 0$ as $t \rightarrow 0$, we obtain from (28)

$$\psi(t) \sim \frac{1}{-\ln(t) + \sqrt{2} + \ln(\sqrt{2} - 1)}, \quad t \rightarrow 0. \quad (30)$$

When $t \rightarrow \infty$, we have $\psi(t) \rightarrow \infty$, and we get from (29)

$$\psi(t) \sim \frac{e}{2}t, \quad t \rightarrow \infty, \quad (31)$$

which agrees with (22).

Since $R(t) = \frac{1}{\psi(t)}$, we have from (28) and (29), after exponentiating

$$e^{-\sqrt{2}} (1 + \sqrt{2}) \frac{R(t) \exp[\sqrt{1 + R^2(t)}]}{1 + \sqrt{1 + R^2(t)}} t = 1, \quad 0 < t \leq 1, \quad (32)$$




Figure 3: A comparison of $R(t)$ (solid curve) and $|A_{500}(t)|^{-\frac{1}{500}}$ (ooo).

and

$$\frac{R(t) \exp \left[\sqrt{1 - R^2(t)} \right]}{1 + \sqrt{1 - R^2(t)}} t = 1, \quad t \geq 1. \quad (33)$$

In Figure 3 we graph the solutions of (32), (33) and the approximate value of $R(t)$ given by $|A_{500}(t)|^{-\frac{1}{500}}$.

Using (2) with $\alpha_n = t^n$, we conclude that the Kapteyn series (7) converges for those $t \in \mathbb{R}$ and $z \in \mathbb{C}$ such that

$$\left| \frac{z \exp \left[\sqrt{1 - z^2} \right]}{1 + \sqrt{1 - z^2}} t \right| < 1. \quad (34)$$

Replacing $z = re^{i\omega}$ in (34), we find that the minimum value of r corresponds to $\omega = \pm \frac{\pi}{2}$. Thus, for $t > 0$, the Kapteyn series (7) will converge inside the circle $|z| < r(t)$, with

$$\frac{r(t) \exp \left[\sqrt{1 + r(t)^2} \right]}{1 + \sqrt{1 + r(t)^2}} t = 1. \quad (35)$$

As it was observed in [8], the circles of convergence of the Kapteyn series (7) and the power series (8) are not equal, the former being slightly smaller than the latter. However, the radius $r(t)$ given by (35) and $R(t)$ are asymptotically equal as $t \rightarrow 0$ and $t \rightarrow \infty$ (see Figure 4).

We summarize our results in the following theorem.

Theorem 2 *The Kapteyn series*

$$F(z, t) = \sum_{n=1}^{\infty} t^n J_n(nz),$$

converges for those $t \in \mathbb{R}$ and $z \in \mathbb{C}$ such that

$$\left| \frac{z \exp \left[\sqrt{1 - z^2} \right]}{1 + \sqrt{1 - z^2}} t \right| < 1.$$

The function $F(z, t)$ admits the power series representation

$$F(z, t) = \sum_{n=1}^{\infty} A_n(t) z^n,$$




Figure 4: A comparison of $r(t)$ (solid curve) and $R(t)$ (ooo).

with coefficients

$$A_n(t) = \frac{(-1)^n}{n!} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n}{k} \left(k - \frac{n}{2}\right)^n t^{n-2k}$$

and radius of convergence $R(t)$ defined by the implicit equations

$$e^{-\sqrt{2}} \left(1 + \sqrt{2}\right) \frac{R(t) \exp \left[\sqrt{1 + R^2(t)} \right]}{1 + \sqrt{1 + R^2(t)}} t = 1, \quad 0 < |t| \leq 1,$$

$$\frac{R(t) \exp \left[\sqrt{1 - R^2(t)} \right]}{1 + \sqrt{1 - R^2(t)}} t = 1, \quad |t| \geq 1.$$

4 The general problem

We shall now consider the problem of finding a_n in terms of α_n , where

$$\sum_{n=1}^{\infty} a_n z^n = \sum_{n=1}^{\infty} \alpha_n J_n(nz), \quad (36)$$

for arbitrary a_n and α_n . Rewriting the power series of $J_n(nz)$ [8]

$$J_n(nz) = \sum_{j=0}^{\infty} \frac{(-1)^j}{j! (n+j)!} \left(\frac{nz}{2}\right)^{n+2j}$$

in the form

$$J_n(nz) = \left(\frac{n}{2}\right)^n \sum_{j=0}^{\infty} \frac{\cos\left(\frac{\pi}{2}j\right) \left(\frac{n}{2}\right)^j}{\left(\frac{j}{2}\right)! \left(n + \frac{j}{2}\right)!} z^{n+j},$$

we have,

$$\begin{aligned} \sum_{n=1}^{\infty} \alpha_n J_n(nz) &= \sum_{n=1}^{\infty} \alpha_n \left(\frac{n}{2}\right)^n \sum_{j=0}^{\infty} \frac{\cos\left(\frac{\pi}{2}j\right) \left(\frac{n}{2}\right)^j}{\left(\frac{j}{2}\right)! \left(n + \frac{j}{2}\right)!} z^{n+j} \\ &= \sum_{k=1}^{\infty} \sum_{n=1}^k \alpha_n \left(\frac{n}{2}\right)^n \frac{\cos\left[\frac{\pi}{2}(k-n)\right] \left(\frac{n}{2}\right)^{k-n}}{\left(\frac{k-n}{2}\right)! \left(\frac{k+n}{2}\right)!} z^k. \end{aligned}$$

Therefore, we obtain from (36) that

$$a_n = \sum_{n=1}^k \alpha_n \frac{\cos \left[\frac{\pi}{2} (k-n) \right] \left(\frac{n}{2} \right)^k}{\left(\frac{k-n}{2} \right)! \left(\frac{k+n}{2} \right)!}$$

or

$$a_n = \sum_{n=1}^k \alpha_n \frac{\cos \left[\frac{\pi}{2} (k-n) \right]}{(k-n)!! (k+n)!!} n^k. \quad (37)$$

In particular, setting $\alpha_n = t^n$ in (37), we recover (17) .

Acknowledgement 3 *We would like to thank our colleague Michael Adams and the anonymous referees, for extremely useful comments on earlier versions of this paper.*

References

- [1] M. Abramowitz and I. A. Stegun. *Handbook of Mathematical Functions*. Dover, New York, 9th ed., 1972.
- [2] F. W. Bessel. Analytische Auflösung der Keplerschen Aufgabe. *Berliner Abh.*, pages 49–55, 1819.
- [3] P. Colwell. *Solving Kepler’s equation over three centuries*. Willmann-Bell Inc., Richmond, VA, 1993.
- [4] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Academic Press Inc., San Diego, CA, sixth edition, 2000.
- [5] W. Kapteyn. Recherches sur les fonctions de Fourier–Bessel. *Ann. Sci. École Norm. Sup.*, x(3):91–120, 1893.
- [6] W. Kapteyn. Over Bessel’sche Functiën. *Nieuw Archief voor Wiskunde*, xx:116–127, 1893.
- [7] J. L. Lagrange. Sur le problème de Kepler. *Hist. de l’Acad. R. des Sci. de Berlin*, xxv:204–233, 1771.
- [8] G. N. Watson. *A treatise on the theory of Bessel functions*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1995.